



Pyrethroid



Capsule
Suspension

PRESCRIPTION TREATMENT[®] brand **CY-KICK[®] ES**

PRESSURIZED

CRACK & CREVICE[®] RESIDUAL

For Use by Commercial Applicators Only

KILLS: Ants, Aphids, Armyworms, Bees, Beetles, Bugs (Boxelder, Lace and Plant) Carpenter Ants, Carpenter Bees, Carpet Beetles, Caterpillars, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Firebrats, Flies, Flour Beetles (Confused, Red, Saw-toothed), Fruitworm, Fungus Gnat Larvae, Grasshoppers, Leafminers, Leafrollers, Loopers, Hornets, Indian Meal Moths, Mediterranean Flour Moths, Millipedes, Mosquitoes, Psylla, Rice Weevils, Scale Insects, Scorpions, Silverfish, Sowbugs, Spiders, Thrips, Ticks, Wasps and Yellowjackets.

FOR USE IN AND AROUND: Apartments, Homes, Hospitals, Hotels, Food/Feed Handling Establishments, Restaurants, Hotels, Hospitals and Nursing Homes (Non-patient Areas), Motels, Hobby Greenhouses, Interiorscapes, Office Buildings, Schools*, Transportation Equipment (Buses, Boats, Ships, Trains, Planes†) and Warehouses and Other Commercial and Industrial Buildings.

* Do not apply to classrooms when in use.

† Do not use in aircraft cabins.

ACTIVE INGREDIENTS:

Cyfluthrin 0.1%

OTHER INGREDIENTS:

..... 99.9%

Contains petroleum distillates.

TOTAL: 100.0%

EPA Reg. No. 499-303 EPA Est. No. 499-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See below for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then

give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment.

NOTE TO PHYSICIAN: This product contains petroleum distillates. Do not induce vomiting. Vomiting may pose an aspiration pneumonia hazard.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas

where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not spray directly into any electronic equipment such as radios, televisions, computers, etc. Do not spray where electrical short circuits might result, such as in wall outlets, conduits, motors, switches, etc. Product should only be used when can temperature is above 60°F. If can temperature is below 60°F, store at room temperature until a temperature above 60°F is reached.



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.



CY-KICK® CS PRESSURIZED Crack & Crevice® Residual

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

SHAKE WELL BEFORE USING

Avoid contamination of food, utensils, and food preparation areas. Keep children and pets off treated areas until spray has dried. Do not use in classrooms while in use. Do not use in rooms while occupied by the elderly or infirm. Reapplications may be made at 10 day intervals.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool area away from heat or open flame.

PESTICIDE DISPOSAL: Waste resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: This container may be recycled in the few but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label (DO NOT PUNCTURE!). If recycling is not available, wrap the container and discard in the trash.

HOW TO USE WHITMIRE MICRO-GEN CRACK AND CREVICE® INJECTION SYSTEM

CRACK AND CREVICE® TREATMENTS: Use CY-KICK® CS PRESSURIZED only with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment. Inject CY-KICK® CS PRESSURIZED into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic area and treat at the rate of five to ten seconds per three cubic feet. Several holes may be required in long running voids.

SPOT TREATMENTS: To make a spot treatment release product onto the surface. Do not exceed 2 sq. ft. for each spot treatment.

INDOOR TREATMENTS

CRAWLING INSECTS - Cockroaches, Spiders, Waterbugs, Silverfish, Crickets, Firebrats: Apply as a spot treatment and/or Crack and Crevice® treatment into hiding places such as cracks and crevices, moist areas, and openings around pipes. Treat floors, walls, under pallets,

behind shelves and drawers. Apply directly on insects in these locations where possible.

TICKS: Thoroughly treat infested and egg laying sites near pet resting areas such as cracks and crevices, behind baseboards, door and window frames.

BOOKLICE & SILVERFISH: Inject into cracks and crevices associated with baseboards, door frames, book cases and other harborage sites.

ANTS: Treat trails, nests and points of entry around doors, book cases and storage areas. Apply directly on insects in these locations where possible.

CARPET BEETLES: Make applications on and under edge of floor coverings, under carpets and furniture and in closets or other localities where these insects are found.

STORED PRODUCT PESTS: Make spot and/or Crack and Crevice® treatment to localized areas including shelves, cupboards, walls and floors where food products suspected of being infested are stored. Remove or cover food before treatment. This treatment is supplemental to good housekeeping, since it will have no effect on insects within packaging. Infested food packaging should be destroyed.

FOOD HANDLING ESTABLISHMENTS

Applications of CY-KICK® CS PRESSURIZED are permitted in both food/feed and non-food/feed areas of food/feed handling establishments as spot treatments or Crack and Crevice® treatments. Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Food/feed areas are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

SPOT AND CRACK AND CREVICE® APPLICATION: Spot or Crack and Crevice® applications may be made while the facility is in operation, except in Federally inspected meat and poultry plants. Food should be covered or removed from area being treated. Do not apply directly to food or food handling surfaces. May be used in federally inspected meat and poultry plants.

INDOOR AND OUTDOOR TREATMENTS

CARPENTER ANTS: Locate source of infestation. Inject 5-10 seconds of product into insect tunnels and cavities. Treat trails, points of entry or exits from voids such as around doors and windows. Treatment of trails on tree trunks or fences is permitted.

FOR ACTIVE AND HIBERNATING STAGES OF BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES AND CLOVER MITES:

Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames.

OUTDOOR TREATMENTS

FLYING INSECTS: Treat outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where insects light or congregate.

Treat normal fly resting areas such as ceilings, upper corners of the area, around windows and doors, light fixtures and other surfaces known to attract flies. Hold can one to two feet from target surface and apply at the rate of one 20 oz. can per 250-375 square feet of surface area. Treat so as to lightly moisten the target surface. For best results initiate treatment at the beginning of the fly season and repeat weekly during periods of heavy fly activity.

CARPENTER BEES: Locate entrance holes. Inject 5 - 10 seconds of product into entrance holes. Seal bee entrances after application.

PERIMETER TREATMENT: For crawling insects, apply CY-KICK® CS PRESSURIZED to cracks and crevices where insects may enter premises. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits, around attic vents, cracks and crevices, openings around windows, doors and pipes. Treatment of infested fences and tree holes is permitted.

To help prevent infestations of buildings treat a band of soil 6 to 10 feet wide around and adjacent to the building. Also, treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. A rate of 14 seconds per 100 square feet is desirable.

CRAWLING INSECTS: For treatment of localized infestations of these insects in outdoor areas which are landscaped, planted, mulched or unmaintained, treat infested areas. For ants, wet hills and runways. Treatment of infested fences and tree holes is permitted.

TREATMENT FOR IN GROUND SERVICE BOXES:

Ants, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Fire Ants, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders (such as Black Widow and Brown Recluse), Springtails. Use actuator or other Whitmire Micro-Gen equipment to treat open boxes and exposed surfaces. Do not contact any electrical wiring or mechanical parts of meter.

A PRESCRIPTION TREATMENT® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682

© 2001 Whitmire Micro-Gen Research Laboratories, Inc.

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.

Revised: 9/01